

ON THE RANGE OF VALIDITY OF THE QCD-IMPROVED PARTON MODEL

A. SZCZUREK

*H. Niewodniczanski Institute of Nuclear Physics, ul. Radzikowskiego 152,
Kraków, Poland*

E-mail: szczurek@alf.ifj.edu.pl

Based on the world DIS data we extract $F_2^p - F_2^n$ as a function of Bjorken x and photon virtuality Q^2 . The standard PDFs fail to describe the experimental data below $Q^2 < 7 \text{ GeV}^2$. The trend of the data is approximately explained by our recent model.

Two methods were used to extract experimental values of $D_G = F_2^p - F_2^n$:

$$\begin{aligned} D_G &= F_2^p(x, Q^2) \cdot \left(1 - \frac{F_2^n(x, Q^2)}{F_2^p(x, Q^2)} \right), \\ D_G &= 2F_2^d(x, Q^2) \cdot \left(\frac{1 - \frac{F_2^n(x, Q^2)}{F_2^p(x, Q^2)}}{1 + \frac{F_2^n(x, Q^2)}{F_2^p(x, Q^2)}} \right). \end{aligned} \quad (1)$$

The two methods above are of course equivalent when nuclear effects are neglected. They are equivalent also in a less obvious case when nuclear effects are taken into account and F_2^n/F_2^p is replaced by $2F_2^d/F_2^p - 1$ ¹. As an example in Fig.1 I show the Q^2 dependence of $F_2^p - F_2^n$ for some selected values of Bjorken x . The results obtained with the two methods agree (compare left and right column). The theoretical results obtained with different recent PDF's³ are shown for comparison. Above $Q^2 > 7 \text{ GeV}^2$ the parton model describes the experimental data well. However, the predictions of the parton model clearly deviate from the experimental data below $Q^2 < 7 \text{ GeV}^2$, especially for Bjorken x between 0.2 and 0.45. The predictions of our recent model² are shown by the thick solid line and reproduce the main trend of the data. Because the VDM contribution to $F_2^p - F_2^n$ almost cancels¹, the experimental data seem to present a real modification of the parton model at small resolution scales. This modification can be viewed formally as higher twist effects¹.




Figure 1: $D_G(Q^2)$ for some selected values of Bjorken x .

References

1. A. Szczurek and V. Uleshchenko, *Phys. Lett. B* **475**, 120 (2000).
2. A. Szczurek and V. Uleshchenko, *Eur.Phys.J. C* **12**, 663 (2000).
3. M. Glück *et al*, *Z. Phys. C* **67**, 433 (1995);
A.D. Martin *et al*, *Eur. Phys. J. C* **4**, 463 (1998);
H.L. Lai *et al*, hep-ph/9903282.